

20-20-20

FOLIAR FEED FOR COTTON

Water Soluble Fertilizer

GUARANTEED ANALYSIS

Total Nitrogen (N)	20.00%
3.90% Ammoniacal Nitrogen	
5.80% Nitrate Nitrogen	
10.30% Urea Nitrogen	
Available Phosphate (P₂O₅)	20.00%
Soluble Potash (K₂O)	20.00%
Boron (B)	0.020%
Copper (Cu).....	0.050%
0.05% Chelated Copper	
Iron (Fe)	0.100%
0.10% Chelated Iron	
Manganese (Mn)	0.050%
0.05% Chelated Manganese	
Molybdenum (Mo).....	0.0005%
Zinc (Zn)	0.050%
0.05% Chelated Zinc	
Chlorine (Cl) Maximum	1.000%

Derived From: Ammonium Phosphate, Potassium Nitrate, Urea, Sodium Borate, Copper EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA, and Sodium Molybdate. Chelating Agent is Ethylenediamine Tetra Acetate (EDTA).

DIRECTIONS FOR USE ON COTTON			
Crop	* Rate	Covers	
Application			
Seedling Cotton	1 lb.	1 Acre Directed Band	2 applications at 7-10 day intervals with normal spraying equipment
Bloom Stage	2 lbs.	1 Acre Directed Band	2 applications at 7-10 day intervals
* Broadcast rate for all stages is 3# per acre.			

CAUTION: THIS PRODUCT CONTAINS BORON, COPPER, IRON, MANGANESE, ZINC, AND MOLYBDENUM. SHOULD BE USED AS RECOMMENDED. MISUSE MAY PROVE HARMFUL. DO NOT USE WITH DINITRO COMPOUNDS, DORMANT OIL, BORDEAUX, SPRAY LIME, OR HIGHLY ALKALINE SPRAY MATERIALS. ALWAYS ADD 20-20-20 TO SPRAY MIXTURE AS THE LAST INGREDIENT.

NET WEIGHT
11.35 KG/25 LBS.